

A Transition Amplitude on the Gauge Orbit Space*

Sergey V. SHABANOV**

*Service de Physique Theorique de Saclay,
CEA-Saclay, Gif-sur-Yvette Cedex, F-91191,
France*

Abstract

A general procedure for deriving the path integral representation of a transition amplitude on the gauge orbit space having a non-trivial topology is proposed. The path integral formula appears to be modified by including trajectories reflected from the physical configuration space boundary into the sum over paths. A solution of the Gribov problem of gauge fixing ambiguities is given in the framework of the path integral modified.

Submitted to "The Proceedings of the II International Workshop on Constrained Systems", Montepulchiano, Italy, June, 1993.

*Work supported by the MRT grant of the government of France.

**On leave from: *Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, P.O.Box 79, Moscow, Russia*

e-mail address: shabanov@amoco.saclay.cea.fr

It is well-known that quantum dynamics strongly depends on a configuration space topology. For example, a spectrum of a free particle moving on a line is continuous, while the same system on a circle has a discrete spectrum. Therefore one can expect that the path integral (PI) representation of a transition amplitude [1],[2] depends on the configuration space topology. Wave functions of systems with topologically non-trivial configuration spaces usually obey some boundary conditions. For instance, wave functions of a particle on a circle satisfy periodic boundary conditions, while wave functions of a particle in a box must vanish at the box walls. Thus, the problem of constructing the PI formalism for such theories is to take into account boundary conditions which contain all information about the configuration space topology.

Naively, one would think that it is sufficient to restrict the integration domain in the path integral to obtain a transition amplitude for a particle in a box or on a circle. However, we know from the operator formalism that transition amplitudes for two systems with the same Hamiltonian and the same volume of the configuration space might be different if they have different boundary conditions. A trivial example is a particle moving in a one-dimensional interval of size L . Zero boundary conditions imply infinite walls attached at the interval ends (a particle in a box), while the periodic ones are for a particle on a circle. The corresponding transition amplitudes are different [3]-[5]. Thus, a restriction of the integration domain does not lead to a correct solution of the problem.

For simplest systems, like a particle in a box or on a circle, a correct PI formula for a transition amplitude is obtained by including trajectories reflected from the box walls or trajectories with all possible winding numbers, respectively, into the Feynman sum over paths [3]- [5] rather than by restricting the integration domain. The method of reflected trajectories can be successfully applied to systems with more complicated configuration spaces [4] (see for a review [6] and references therein). This structure of PI is established even for such unusual systems as the q -deformed ones [7].

Gauge theories (or the first class-constrained systems [8]) give another example of this kind. Their main feature is the existence of unphysical variables whose evolution is not determined by equations of motion and appears to be completely arbitrary. This arbitrariness occurs through the gauge invariance of a corresponding Lagrangian [8]. In the whole (total) configuration (or phase) space, there are points related to each other by gauge transformations depending on arbitrary functions of time. These points form orbits of the gauge group. Any motion of a system along a gauge orbit does not lead to any change of a physical state of a system [8]. Two physically distinct states correspond to different gauge orbits. Therefore, to construct a physical configuration space CS_{ph} (each its point corresponds, by definition, to just one physical state of a system), one should stick together all points of each gauge orbit. In the mathematical language, the physical configuration space is the quotient of the total configuration space (CS) by the gauge group G ,

$$CS_{ph} = CS/G . \quad (1)$$

The gauge orbit space (1) might differ from an ordinary Euclidean space [9]-[11] and possess a non-trivial topology even if the total CS is assumed to be Euclidean. For example, it turns out to be compact for QCD_2 [12],[13] and coincides with the Weyl cell (being a polyhedron) for a compact gauge group [12].

In the present paper, we shall give a general recipe for deriving the PI representation of a transition amplitude on the gauge orbit space (1).

Consider a quantum theory determined by the Schroedinger equation

$$\left(-\frac{1}{2}\left\langle\frac{\partial}{\partial x},\frac{\partial}{\partial x}\right\rangle+V(x)\right)\psi_E=E\psi_E. \quad (2)$$

The eigen-functions ψ_E are normalized by the condition

$$\int_{\mathbb{R}^N} dx \psi_E^*(x) \psi_{E'}(x) = \delta_{EE'}. \quad (3)$$

We assume x to realize a linear representation of a compact group G so that its action is given by a linear operator $x \rightarrow T(\omega)x$, and $V(Tx) = V(x)$; $\langle x, y \rangle = \sum_1^N x_i y_i = \langle Tx, Ty \rangle$ is an invariant scalar product in the representation space that is isomorphic to $\mathbb{R}^N$. The theory turns into the gauge one if we require that physical states are annihilated by operators generating G -transformations of x , $\hat{\sigma}_a \Phi(x) = 0$. These conditions determine a physical subspace in the Hilbert space. By definition, we have $\exp(\omega_a \hat{\sigma}_a) \psi(x) = \psi(T(\omega)x)$. Therefore, the physical states are G -invariant

$$\Phi(T(\omega)x) = \Phi(x), \quad (4)$$

where ω runs over the group manifold.

Let a number of physical degrees of freedom in the system is equal to M , then a number of independent constraints is $N - M$. Suppose we would like to span the physical configuration space $K \sim \mathbb{R}^N/G$ by coordinates ranging a gauge condition surface $F(x) = 0$. We assume the gauge condition to be complete, meaning that there is no unphysical degree of freedom left. Let $u \in \mathbb{R}^M$ be a parameter of the gauge condition surface $x = f(u)$ such that $F(f(u))$ identically vanishes for all $u \in \mathbb{R}^M$. The gauge condition surface must cross each gauge orbit at least once so that u could serve as a coordinate spanning K . In other words, any particular configuration x can be transferred onto the gauge condition surface by a gauge transformation, i.e. x can be represented as follows

$$x = x(\theta, u) = T(\theta)f(u). \quad (5)$$

Physical states (4) are independent of the variables θ spanning the gauge group manifold. Therefore, a projection of the Hamiltonian operator on the physical subspace can be done by introducing the curvilinear coordinates (5) in the Laplace operator entering into (2) and by dropping all terms containing derivatives with respect to θ in it. Provided each gauge orbit intersects a gauge condition surface just once, we would have a quantum theory on a curved configuration space which is topologically equivalent to $\mathbb{R}^M$. A construction of the PI representation for a transition amplitude in this case does not give rise to any serious problem, and is known [5], [6]. Troubles appear when a gauge condition does not completely fix a gauge arbitrariness, meaning that there are configurations on the gauge condition surface which are connected with each other by gauge transformations. The residual gauge arbitrariness cannot decrease a number of physical degrees of freedom, but it does reduce their configuration space.

The existence of residual gauge transformations means that the gauge condition surface intersects each gauge orbit not only once. All the intersection points are, obviously, gauge copies of one of them [14]. The latter may occur due to two reasons. The first one is that a gauge condition looks preferable because of physical reasons in spite of its ambiguity. Another reason is hidden in a mathematical structure of gauge systems. One cannot always fix a gauge without any ambiguity due to a non-trivial topology of the gauge orbit space. Yang-Mills fields give an example of such a kind [14],[15].

As has been pointed out above, a configuration space topology is taken into account by boundary conditions imposed on wave functions. Therefore, to construct the PI representation of a transition amplitude on the gauge orbit space, one should first find boundary conditions for physical states appearing upon projecting the Hamiltonian operator on the orbit space, and then to derive PI corresponding to them. To go over our program, we shall introduce the curvilinear coordinates (5) to remove unphysical degrees of freedom. The boundary conditions determining the gauge orbit space topology will be shown to result from analytical properties of physical wave functions.

The metric tensor in the new coordinates reads

$$\langle dx, dx \rangle = \langle df, df \rangle + 2\langle df, d\theta f \rangle + \langle d\theta f, d\theta f \rangle \equiv g_{AB} dy^A dy^B, \quad (6)$$

where we have used the G -invariance of the bilinear form $\langle, \rangle$ and put $d\theta = T^{-1}dT$ and $dy^1 \equiv du$, $dy^2 \equiv d\theta$. Therefore,

$$\int_{\mathbb{R}^N} dx = \int_G \wedge d\theta \int_K d^M u \mu(u); \quad (7)$$

here $\mu(u) = (\det g_{AB})^{1/2}$, K is a subdomain in $\mathbb{R}^M$ such that the mapping (5), $K \otimes G \rightarrow \mathbb{R}^N$, is one-to-one. To determine K , one should find transformations $\theta, u \rightarrow \hat{s}\theta, \hat{s}u$, $\hat{s} \in \tilde{S}_F$, which leave x untouched, $x(\hat{s}\theta, \hat{s}u) = x(\theta, u)$. Obviously, $\tilde{S}_F = T_e \times S_F$ where T_e is a group of translations of θ through periods of the group manifold G and $\hat{s}u = u$, $\hat{s} \in T_e$, while the set S_F is obtained by solving the following equation

$$F(T_s f(u)) = 0 \quad (8)$$

with respect to a gauge transformation operator T_s . Indeed, assuming Eq.(8) to have non-trivial solutions (the trivial one $T_s = 1$ always exists by the definition of $f(u)$) we observe that all points $T_s f$ belong to the gauge condition surface and, hence, $T_s f(u) = f(u_s)$, $u_s = u_s(u)$. Consider transformations of θ generated by the group shift $T(\theta) \rightarrow T(\theta)T_s^{-1} = T(\theta_s)$, $\theta_s = \theta_s(\theta, u)$. Setting $\hat{s}u = u_s$ and $\hat{s}\theta = \theta_s$ we see that the transformations $\hat{s} \in S_F$ leave $x = x(\theta, u)$ untouched. To avoid a "double" counting in the scalar product integral (7) (cf. (3)), one has to restrict the integration domain for u to the quotient $\mathbb{R}^M/S_F = K$. The modular domain K can also be determined by the requirement that a part of the gauge condition surface $x = f(u)$, $u \in K \subset \mathbb{R}^M$ has just one common point with any gauge orbit.

A choice of the modular domain parametrization is not unique. Suppose we find a part of a given gauge condition surface such that each gauge orbit is represented by one point on it, i.e. we identify K with a concrete subregion in $\mathbb{R}^M$. Apparently, any subregion

among $\hat{s}K = K_s$, $\hat{s} \in S_F$, can serve as the integration region in (7). Having chosen a concrete parametrization (coordinates) of K , we fix a representation of S_F by functions $\hat{s}u = u_s(u)$, $u \in K$, $u_s \in K_s$, $K_s \cap K_{s'} = \emptyset$ for any $\hat{s} \neq \hat{s}'$ and $\mathbb{R}^M = \cup_s K_s$ up to a set of zero measure being a unification of the boundaries ∂K_s .

The functions $u_s(u)$ might not have an analytical continuation to the whole covering space $u \in \mathbb{R}^M$. Let u_s and $u_{s'}$ have an analytical continuation to $\mathbb{R}^M$. The latter would mean that their composition $u_s \circ u_{s'}$ exists and must determine an operator $T_{ss'}$ obeying (8) since there should be $T_{ss'} = T_s T_{s'}$. However, a composition of two solutions to (8) does not always give a new solution because $T_s = T_s(u)$ and $T_{s'} = T_{s'}(u)$ are functions of u , i.e. in a general case $F(T_s(u)T_{s'}(u)f(u)) = F(T_s(u)f(u_{s'})) \neq 0$ since $u_{s'} \neq u$, whereas $F(T_s(u)f(u)) = 0$. Herein we restrict ourselves just by pointing out this fact and by illustrating it with a simple example (see it below). We shall assume that the functions u_s are analytical only on K , while u_s^{-1} maps K_s on K , which is sufficient for what follows. So, we admit the set of transformations S_F to be not a group since a composition of its elements might not be uniquely defined. Notice also that residual gauge transformations may not form a group for Yang-Mills theories [16].

We define an orientation of K_s so that for all $\hat{s} \in S_F$, $\int_{K_s} du \phi \geq 0$ if $\phi \geq 0$, which provides the following rules

$$\int_{\mathbb{R}^M} d^M u = \sum_{S_F} \int_{K_s} d^M u, \quad (9)$$

$$\int_K d^M u |J_s(u)| = \int_{K_s} d^M u, \quad (10)$$

where $J_s(u) = Du_s/Du$ is the Jacobian, the absolute value of J_s has been inserted into the left-hand side of (10) for preserving the positive orientation of the integration domain. We assume the Jacobian μ in (7) to be positive on the domain K chosen, otherwise one should take its absolute value in accordance with our orientation rules.

Remark. A number of elements in S_F can depend on u . We define a region $\mathbb{R}_\alpha^M \subseteq \mathbb{R}^M$ such that $S_F = S_\alpha$ has a fixed number of elements for all $u \in \mathbb{R}_\alpha^M$. Then $K = \cup_\alpha K_\alpha$, $K_\alpha = \mathbb{R}_\alpha^M / S_\alpha$, $\mathbb{R}^M = \cup_\alpha \mathbb{R}_\alpha^M$. The sum in (9) implies $\sum_{S_F} = \sum_\alpha \sum_{S_\alpha}$ and K_s in (9-10) carries an additional suffix α . In what follows we shall omit it and use the simplified notations (9-10) to avoid complications of formulas. The suffix α can be easily restored by means of the rule proposed above.

Example. Let the whole configuration space be a plane $\mathbb{R}^2$ and the gauge group be an $SO(2)$ -rotation of $\mathbb{R}^2$. Gauge orbits are concentric circles. Any gauge condition $F(\mathbf{x}) = 0$, $\mathbf{x} \in \mathbb{R}^2$, determine a curve $\mathbf{x} = \mathbf{f}(u)$, $u \in \mathbb{R}$, which goes through the origin to infinity (to provide crossing each orbit at least once). So, $T(\theta) = \exp(i\theta\sigma_2)$, σ_2 the Pauli matrix, and $x, f \rightarrow \mathbf{x}, \mathbf{f} \in \mathbb{R}^2$ in (5). Set $f_1 = -u_0$, $f_2 = -\gamma(2u_0 + u)$ for $u < -u_0$ and $f_1 = u$, $f_2 = \gamma u$ for $u > -u_0$ where γ and u_0 are positive constants. The curve $x_{1,2} = f_{1,2}(u)$ touches circles (gauge orbits) of radii $r = u_0$ and $r = u_0\gamma_0$, $\gamma_0 = \sqrt{1 + \gamma^2}$. It intersects twice all circles with radii $r < u_0$ and $r > u_0\gamma_0$, whereas any circle with a radius from the interval $r \in (u_0, u_0\gamma_0)$ has four common points with the gauge condition curve. Therefore, S_F has one nontrivial element for $u \in \mathbb{R}_1 \cup \mathbb{R}_3$, $\mathbb{R}_1 = (-u_0/\gamma_0, u_0/\gamma_0)$, $\mathbb{R}_3 = (-\infty, -3u_0) \cup (u_0, \infty)$ and three nontrivial elements for $u \in \mathbb{R}_2 = (-3u_0, -u_0/\gamma_0) \cup$

$(u_0/\gamma_0, u_0)$. Since points $\mathbf{f}(u_s)$ and $\mathbf{f}(u)$ belong to the same circle (gauge orbit), the functions u_s have to obey the following equation

$$\mathbf{f}^2(u_s) = \mathbf{f}^2(u) . \quad (11)$$

Denoting $S_F = S_\alpha$ for $u \in \mathbb{R}_\alpha$, $\alpha = 1, 2, 3$ (see *Remark* above), we have $S_1 = \mathbb{Z}_2$, $u_s(u) = -u$; S_2 is determined by the following mappings of the interval $K_2 = (u_0/\gamma_0, u_0)$

$$u_{s_1}(u) = -u , \quad u_{s_1} : K_2 \rightarrow (-u_0, -u_0/\gamma_0) ; \quad (12)$$

$$u_{s_2}(u) = -2u_0 + \gamma_0(u^2 - u_0^2/\gamma_0^2)^{1/2}/\gamma , \quad u_{s_2} : K_2 \rightarrow (-u_0, -2u_0) ; \quad (13)$$

$$u_{s_3}(u) = -2u_0 - \gamma_0(u^2 - u_0^2/\gamma_0^2)^{1/2}/\gamma , \quad u_{s_3} : K_2 \rightarrow (-2u_0, -3u_0) ; \quad (14)$$

and for S_3 we get

$$u_s(u) = -2u_0 - \gamma_0(u^2 - u_0^2/\gamma_0^2)^{1/2}/\gamma : (u_0, \infty) \rightarrow (-\infty, -3u_0) . \quad (15)$$

The functions (12-14) do not have a unique analytical continuation to the whole domain $\mathbb{R}_2$ and, hence, their composition is ill-defined. The mappings (12-14) do not form a group. Since they realize a representation of S_α , S_α is not a group.

The physical configuration space is, obviously, isomorphic to $K = \cup K_\alpha$, $K_\alpha = \mathbb{R}_\alpha/S_\alpha$, i.e. K_α is a fundamental domain of $\mathbb{R}_\alpha$ with respect to the action of $S_F = S_\alpha$ in $\mathbb{R}_\alpha$, $\mathbb{R}_\alpha = \cup \hat{s} K_\alpha$, $\hat{s}$ ranges over S_α . Upon solving (8) (or (11)) we have to choose a particular interval as the fundamental domain. We have put $K_2 = (u_0/\gamma_0, u_0)$ in (12-14). Another choice would lead to another form of the functions u_s (to another representation of S_F in $\mathbb{R}_2$). Setting, for example, $K_2 = (-2u_0, -u_0)$ we obtain from (11)

$$u_{s_1}(u) = -4u_0 - u , \quad u_{s_1} : K_2 \rightarrow (-3u_0, -2u_0) ; \quad (16)$$

$$u_{s_2}(u) = -(u_0^2 + \gamma^2(2u_0 + u)^2)^{1/2}/\gamma_0 , \quad u_{s_2} : K_2 \rightarrow (-u_0, -u_0/\gamma_0) ; \quad (17)$$

$$u_{s_3}(u) = (u_0^2 + \gamma^2(2u_0 + u)^2)^{1/2}/\gamma_0 , \quad u_{s_3} : K_2 \rightarrow (u_0/\gamma_0, u_0) . \quad (18)$$

To find group elements $T_s(u)$ corresponding to $u_s(u)$, one should solve the equation $T_s \mathbf{f}(u) = \mathbf{f}(u_s)$. Setting $T_s = \exp(i\omega_s \sigma_2)$ and taking u_s from (12-14) we find

$$\omega_{s_1}(u) = \pi ; \quad (19)$$

$$\omega_{s_2}(u) = \frac{3\pi}{2} - \sin^{-1} \left(\frac{u_0}{\gamma_0 u} \right) - \tan^{-1} \gamma ; \quad (20)$$

$$\omega_{s_3}(u) = \frac{\pi}{2} + \sin^{-1} \left(\frac{u_0}{\gamma_0 u} \right) - \tan^{-1} \gamma , \quad (21)$$

where $u \in K_2 = (u_0/\gamma_0, u_0)$. Elements of $S_{1,3}$ are obtained analogously. It is readily seen that $\Omega_{s_1} \Omega_{s_2} \neq \Omega_{s_3}$, etc., i.e. the elements Ω_s do not form a group. An alternative choice of K_2 results in a modification of the functions (19-21).

For the curvilinear coordinates (5) we have $\det g_{AB} = (\mathbf{f}', \mathbf{f})^2 = \mu^2(u)$. Set $K = \cup_\alpha K_\alpha$, $K_1 = (0, u_0/\gamma_0)$, $K_2 = (u_0/\gamma_0, u_0)$, $K_3 = (u_0, \infty)$, i.e. $K = \mathbb{R}_+$, then $\int_{-\infty}^{\infty} du = \sum_\alpha \int_{\mathbf{R}_\alpha} du$ and (9) means that the upper integral limit is always greater than the lower one, for example,

$$\int_{\mathbf{R}_2} du = \left(\int_{-3u_0}^{-2u_0} + \int_{-2u_0}^{-u_0} + \int_{-u_0}^{-u_0/\gamma_0} + \int_{u_0/\gamma_0}^{u_0} \right) du ,$$

where the terms of the sum correspond to integrations over $\hat{s}_3 K_2$, $\hat{s}_2 K_2$, $\hat{s}_1 K_2$ and K_2 , respectively (cf. (12-14)). The following chain of equalities is to illustrate the rule (10)

$$\int_{\hat{s}_3 K_2} du_{s_3} = \int_{-3u_0}^{-2u_0} du_{s_3} = \int_{u_0}^{u_0/\gamma_0} du J_{s_3} = - \int_{u_0/\gamma_0}^{u_0} du J_{s_3} = \int_{K_2} du |J_{s_3}| ; \quad (22)$$

the last equality results from $J_{s_3} = du_{s_3}/du < 0$ (cf. (14)).

By means of the curvilinear coordinates (5) we can naturally incorporate a gauge condition chosen into the Dirac operator method [8] of quantizing first-class constrained systems. Solutions of the equation $\hat{\sigma}_a \tilde{\Phi}(x) = 0$ are given by functions independent of θ ,

$$\tilde{\Phi}(x) = \tilde{\Phi}(T(\theta)f(u)) = \tilde{\Phi}(f(u)) = \Phi(u) , \quad (23)$$

because $\hat{\sigma}_a$ generate shifts of θ and leave u untouched. To obtain a physical Hamiltonian, one has to write the Laplacian in (2) via the new variables (5) and omit all terms containing derivatives with respect to θ . In so doing, we get

$$\hat{H}_{ph}^f \Phi_E(u) = \left(\frac{1}{2} \hat{p}_i g_{ph}^{ij} \hat{p}_j + V_q(u) + V(f(u)) \right) \Phi_E(u) = E \Phi(u) ; \quad (24)$$

here we have introduced hermitian momenta $\hat{p}_i = -i\mu^{-1/2} \partial_j \circ \mu^{1/2}$, $\partial_j = \partial/\partial u^j$; the metric g_{ph}^{ij} in the physical configuration space is the 11-component of a tensor g^{AB} inverse to g_{AB} , $g^{AC} g_{CB} = \delta_B^A$, $g_{ph}^{ij} = (g^{11})^{ij}$, $i, j = 1, 2, \dots, M$; a quantum potential

$$V_q = \frac{1}{2\sqrt{\mu}} (\partial_i g_{ph}^{ij}) \partial_j \sqrt{\mu} + \frac{1}{2\mu} g^{ij} \partial_i \partial_j \sqrt{\mu} \quad (25)$$

appears due to the chosen ordering of the operators $\hat{u}^i$ and $\hat{p}_i$ in the Laplace-Beltrami operator. Because of (23), the scalar product (3) is reduced to

$$\int_{\mathbf{R}^N} dx \Phi_E^*(u) \Phi_{E'}(u) \rightarrow \int_K d^M u \mu(u) \Phi_E^*(u) \Phi_{E'}(u) = \delta_{EE'} , \quad (26)$$

where a gauge orbit volume (integral over G (see (7)) has been included into norms of physical states, which we denoted by the arrow in (26). A construction of an operator description of a gauge theory in a given gauge condition is completed.

Notice, in this approach the variables u appear to be gauge-invariant, they parametrize the physical configuration space $CS_{ph} = K = \mathbf{R}^N/G$. Two different choices of $f(u)$ in (5) implies two different parametrizations of CS_{ph} related to each other by a change of variables $u = u(\tilde{u})$ in (24-26). Therefore quantum theories with different f 's are unitary equivalent [18], [6].

To illustrate this statement, we consider again the simplest case [17] $G = SO(2)$, $M = 1$, $g_{ph} = \mathbf{f}^2(u)/\mu^2(u)$, and compare descriptions in the coordinates (5) and in the polar ones ($f_1 = r, f_2 = 0$). With this purpose we change variables $r = r(u) = \sqrt{\mathbf{f}^2(u)}$ in (24-26). For $u \in K$ the function $r(u)$ is invertible, $u = u(r), r \in \mathbf{R}_+$. Simple straightforward calculations [17] lead us to the following equalities $\hat{H}_{ph}^f = 1/2 \hat{p}_r^2 + V_q(r) + V$, $\hat{p}_r =$

$-ir^{-1/2}\partial_r \circ r^{1/2}$, $V_q = -(8r^2)^{-1}$, $\int_K du\mu = \int_0^\infty dr r$. It is nothing but quantum mechanics of a radial motion on a plane. All theories with different f 's are unitary equivalent to it and, therefore, to each other. One should stress that the operator ordering we obtained by applying the Dirac method plays the crucial role in providing this unitary equivalence. Another ordering of operators in (24) would break this property.

A few observations resulting from our consideration have to be emphasized.

1. All regular solutions of (24) have a unique analytical continuation to the whole space $u \in \mathbb{R}^M$, and they are S_F -invariant,

$$\Phi_E(u_s(u)) = \Phi_E(u) , \quad u \in K . \quad (27)$$

For a proof, we point out that any regular solution of (24) is a projection of a regular G -invariant solution of (2) on K determined by (23). The last equality in (23) defines the analytical continuation of $\Phi_E(u)$ because we assume $f(u)$ to be analytical on $\mathbb{R}^M$; (27) follows from the second equality in (23) and the definition $T_s f(u) = f(u_s)$. Another proof of (27) is to use averaging over the group manifold. Let $\Psi_E(x)$ be a solution to (2). Then a solution to (24) is obtained by averaging $\Psi_E(x)$ over the group manifold. Hence,

$$\Phi_E(u) = \frac{1}{V_G} \int_G d\mu_G(\theta) \Psi_E(T(\theta)f(u)) = \frac{1}{V_G} \int_G d\mu_G(\theta) \Psi_E(T(\theta)T_s f(u)) = \Phi_E(u_s) ,$$

where V_G is a group volume, $d\mu_G$ in the right-invariant Haar measure, $d\mu(\theta_s) = d\mu(\theta)$ with $T(\theta_s) = T(\theta)T_s^{-1}$.

The condition (27) determines the boundary condition which should be taken into account in the PI representation of the transition amplitude on the gauge orbit space.

2. Any amplitude, i.e. the scalar product (26) of two S_F -invariant states, is independent of a CS_{ph} parametrization (of a gauge choice). An S_F -invariant regular function of $u \in \mathbb{R}^M$ is a linear combination of the basis states $\Phi_E(u)$. Our statement follows from the unitary equivalence of the theories (24-26) corresponding different parametrizations of CS_{ph} .

3. The physical Hamiltonian in (24) is S_F -invariant

$$\hat{H}_{ph}^f(u_s) = \hat{H}_{ph}^f(u) , \quad u \in K. \quad (28)$$

Let us write the Laplace-Beltrami operator $\langle \partial/\partial x, \partial/\partial x \rangle = \Delta(\theta, u)$ in the variables (5), push all derivatives ∂_θ in it to the right by commuting them with θ and then set $\partial_\theta = 0$. We denote the operator thus obtained $\Delta_{ph} = \Delta(\theta, u)|_{\partial_\theta=0}$. For any physical state $\Phi = \Phi(u)$, we have $\Delta(\theta, u)\Phi = \Delta_{ph}\Phi$ because $\hat{\sigma} \sim \partial_\theta$. Due to the gauge-invariance, the Hamiltonian in (2) commutes with the constraints, $[\hat{H}, \hat{\sigma}] = 0$ and, hence, $[\Delta(\theta, u), \hat{\sigma}] = 0$ (the potential V is G -invariant). Gathering the definition of Δ_{ph} and G -invariance of Δ we conclude that $\Delta_{ph} = \Delta_{ph}(u)$ is independent of θ (otherwise we would arrive to the contradiction $0 = [\Delta, \hat{\sigma}]\Phi = \hat{\sigma}\Delta\Phi = \hat{\sigma}\Delta_{ph}\Phi \sim \partial_\theta\Delta_{ph}\Phi \neq 0$). Consider now the change of variables $\theta, u \rightarrow \theta_s, u_s$. By its definition $\Delta(\theta_s, u_s) = \Delta(\theta, u)$ and $\partial_\theta \sim \partial_{\theta_s}$ (i.e. ∂_θ does not contain a term proportional ∂_{u_s} since $\partial u_s / \partial \theta = 0$). This yields $\Delta_{ph}(u_s) = \Delta(\theta_s, u_s)|_{\partial_{\theta_s}=0} = \Delta(\theta_s, u_s)|_{\partial_\theta=0} = \Delta(\theta, u)|_{\partial_\theta=0} = \Delta_{ph}(u)$, which completes the proof of (28).

To derive a path integral representation of the quantum theory (24-26), we consider a slice approximation of the transition amplitude $U_t^{ph}(u, u') = \langle u | \exp(-i\hat{H}_{ph}t) | u' \rangle$,

$$U_t^{ph}(u, u') = \lim_{\epsilon \rightarrow 0} \int_K \prod_{k=0}^n (d^M u_k \mu(u_k)) U_\epsilon^{ph}(u, u_n) U_\epsilon^{ph}(u_n, u_{n-1}) \cdots U_\epsilon^{ph}(u_1, u') , \quad (29)$$

where $(n+1)\epsilon = t$, the limit is taken so that $n \rightarrow \infty$, $\epsilon \rightarrow 0$, while t is kept fixed; the infinitesimal evolution operator kernel reads

$$U_\epsilon^{ph}(u, u') = [1 - i\epsilon \hat{H}_{ph}(u)] \langle u | u' \rangle + O(\epsilon^2) . \quad (30)$$

A naive limit in (29) gives a formal PI with a restricted integration domain $K \subset \mathbb{R}^M$. A calculation of such a PI meets difficulties because even a finite dimensional Gaussian integral cannot be explicitly done over a part of an Euclidean space. In addition, a restriction of the PI integration domain is meaningless for systems with boundary conditions appearing due to a non-trivial topology of a configuration space like for a particle in a box or on a circle [3]-[6]. Topological properties of a configuration space are taken into account in PI by including additional “reflected” trajectories into the sum over paths [4],[6] rather than by restricting the PI integration domain. Technically, a relation between transition amplitudes U_t and U_t^{eff} for the same systems (the same Hamiltonian) in a topologically non-trivial CS and in $CS = \mathbb{R}^M$, respectively, is established by means of an operator $\hat{Q}$ containing all information about a CS topology, $\hat{U}_t = \hat{U}_t^{eff} \hat{Q}$ [4],[6]. The same form of PI turns out to be valid for many particular gauge models [6],[10]-[12], [18]. Bellow we shall prove this for our general case.. For deriving a PI formula we shall use a method of an analytical continuation of the unit operator kernel [18],[6].

The unit operator kernel $\langle u | u' \rangle$ has a natural analytical continuation to the unphysical domain $u \in \mathbb{R}^M$. Indeed, due to the S_F -invariance of the basis states (27) and the scalar product form (26) we have

$$\langle u | u' \rangle = \sum_E \Phi_E(u) \Phi_E^*(u') = \sum_{S_F} (\mu(u) \mu(u'))^{-1/2} \delta^M(u - \hat{s}u') , \quad (31)$$

where $u \in \mathbb{R}^M$, $u' \in K$. Representing δ -functions in (31) through the Fourier integral, substituting it into (30) and calculating the action of $\hat{H}_{ph}(u)$ on the unit operator kernel, we obtain

$$U_\epsilon^{ph}(u, u') = \int_{\mathbb{R}^M} \frac{d^M u''}{(\mu \mu'')^{1/2}} U_\epsilon^{eff}(u, u'') Q(u'', u') , \quad (32)$$

$$U_\epsilon^{eff}(u, u'') = \int_{\mathbb{R}^M} \frac{d^M p}{(2\pi)^M} \exp \left[i\epsilon \left(p_j \frac{\Delta_j}{\epsilon} - H^{eff}(u, p) \right) \right] , \quad (33)$$

$$H^{eff}(u, p) = \frac{1}{2} g_{ph}^{ij}(u) p_i p_j + \frac{i}{2} \partial_i g_{ph}^{ij}(u) p_j + V_q(u) + V , \quad (34)$$

$$Q(u'', u') = \sum_{S_F} \delta^M(u'' - \hat{s}u') , \quad u'' \in \mathbb{R}^M , \quad u' \in K , \quad (35)$$

where $\mu'' = \mu(u'')$ and $\Delta_j = u_j - u_j''$. So, the infinitesimal evolution operator kernel (32) has the desired form $\hat{U}_\epsilon^{ph} = \hat{U}_\epsilon^{eff} \hat{Q}$. A next step is to prove the convolution formula

$$U_{2\epsilon}^{ph}(u, u') = \int_K d^M u_1 \mu(u_1) U_\epsilon^{ph}(u, u_1) U_\epsilon^{ph}(u_1, u') \quad (36)$$

$$= \int_{\mathbf{R}^M} \frac{d^M u''}{(\mu \mu'')^{1/2}} U_{2\epsilon}^{eff}(u, u'') Q(u'', u') , \quad (37)$$

$$U_{2\epsilon}^{eff}(u, u'') = \int_{\mathbf{R}^M} d^M u'' U_\epsilon^{eff}(u, u'') U_\epsilon^{eff}(u'', u') , \quad (38)$$

or in the operator form

$$\hat{U}_{2\epsilon}^{ph} = \hat{U}_\epsilon^{eff} \hat{Q} \hat{U}_\epsilon^{eff} \hat{Q} = \hat{U}_{2\epsilon}^{eff} \hat{Q} . \quad (39)$$

The proof is given by the following chain of equalities

$$U_{2\epsilon}^{ph}(u, u') = \sum_{S_F} \int_K d^M u_1 \frac{\mu_1}{(\mu \mu(\hat{s}u_1))^{1/2}} U_\epsilon^{eff}(u, \hat{s}u_1) U_\epsilon^{ph}(u_1, u') = \quad (40)$$

$$= \sum_{S_F} \int_K d^M u_1 |J_s(u_1)|^{1/2} \left(\frac{\mu_1}{\mu} \right)^{1/2} U_\epsilon^{eff}(u, \hat{s}u_1) U_\epsilon^{ph}(\hat{s}u_1, u') = \quad (41)$$

$$= \int_{\mathbf{R}^M} \frac{d^M u''}{(\mu \mu'')^{1/2}} \sum_{S_F} \int_K d^M u_1 |J_s(u_1)| U_\epsilon^{eff}(u, \hat{s}u_1) U_\epsilon^{eff}(\hat{s}u_1, u'') Q(u'', u') , \quad (42)$$

where $\hat{s}u_1 = u_s(u_1)$, $\mu_1 = \mu(u_1)$, $\mu = \mu(u)$ and $\mu'' = \mu(u'')$. To obtain (40), we substitute (32) into (36) and do the integral over u'' . For the transformation (41) we use the relation $\mu(u_s) = \mu(u)/J_s(u)$ (which follows from the S_F -invariance of the measure $d^N x = (\wedge d\theta) d^M u \mu(u) = (\wedge d\theta) d^M u_s \mu(u_s)$) and the S_F -invariance of the kernel (32) (or (30)). The latter results from the obvious relation $\langle \hat{s}u | u' \rangle = \langle u | u' \rangle$, $u \in K$ (cf. (31)) and (28). Equality (42) is derived by substituting (32) into (41) and using the relation $\mu(\hat{s}u_1) = \mu(u)/J_s(u)$ again. Finally, (42) turns into (37) after changing variables $u_1 \rightarrow \hat{s}u_1$ in each term of the sum in (42) by means of the rules (9) and (10).

For a finite time we get (see (39))

$$\hat{U}_t^{ph} = \lim_{\epsilon \rightarrow 0} \hat{U}_\epsilon^{eff} \hat{Q} \hat{U}_\epsilon^{eff} \hat{Q} \dots \hat{U}_\epsilon^{eff} \hat{Q} = \lim_{\epsilon \rightarrow 0} \hat{U}_\epsilon^{eff} \dots \hat{U}_\epsilon^{eff} \hat{Q} = \hat{U}_t^{eff} \hat{Q} , \quad (43)$$

$$U_t^{eff}(u, u'') = \lim_{\epsilon \rightarrow 0} \int_{\mathbf{R}^M} \left(\prod_{k=1}^n d^M u_k \right) U_\epsilon^{eff}(u, u_n) U_\epsilon^{eff}(u_n, u_{n-1}) \dots U_\epsilon^{eff}(u_1, u'') . \quad (44)$$

Formulas (43-44) and (33) with $\Delta_k/\epsilon = \dot{u}_k + O(\epsilon)$ solve the problem of the PI construction. Equalities (44) and (33) determine a standard slice approximation of PI over an Euclidean phase space. Removing the slice regularization in (44) we obtain the path integral

$$U_t^{eff}(u, u'') = \int_{\mathbf{R}^{2M}} \left(\prod_{\tau=0}^t \frac{d^M p(\tau) d^M u(\tau)}{(2\pi)^M} \right) \exp i \int_0^t d\tau (p_j \dot{u}_j - H^{eff}) , \quad (45)$$

where the measure implies a sum over all trajectories $u(\tau)$ going from the initial point $u'' = u(0)$ to the final one $u = u(t)$. The physical transition amplitude is given by (32) ($\epsilon \rightarrow t$) and implies a sum over trajectories going from a *few* initial points $u_s(u') = \hat{s}u'$, $u' = u(0)$, $\hat{s} \in S_F$, to the final one. A trajectory going from one of these points, say, $\hat{s}u' \in K_s$, $\hat{s} \neq 1$, to $u = u(t) \in K$ must cross the boundary ∂K at a point $\tilde{u} = u(\tilde{\tau})$. Suppose for simplicity that $u(\tau) \in K$ if $\tau \in (\tilde{\tau}, t)$ and $u(\tau) \in K_s$ if $\tau \in (0, \tilde{\tau})$. Consider a reflected trajectory composed of two pieces $\hat{s}^{-1}u(\tau)$, $\tau \in (0, \tilde{\tau})$ and $u(\tau)$, $\tau \in (\tilde{\tau}, t)$, i.e. it connects the initial point $u' \in K$, the "reflection" point $\tilde{u} \in \partial K$ and the final point $u \in K$. Due to the S_F -invariance of the effective action, the reflected trajectory gives the same contribution into the sum over paths as the "straight" one $u_s(u') \rightarrow u$. *Therefore the PI modification (43) due to a non-trivial topology of PS_{ph} (or CS_{ph}) means that in addition to "straight" trajectories $u' \rightarrow u$, the reflected trajectories $u' \rightarrow \partial K \rightarrow u$ must be included into the sum over paths.*

Contributions of the "reflected" trajectories modify a semiclassical approximation of PI [6]. For instance, in minisuperspace cosmological models with gauge fields [19], the Hartley-Howking wave function of the universe depends on the structure of a physical configuration (phase) space of gauge fields [20].

An analysis of quantum Green functions on the gauge orbit space can be found in [18].

Applications to gauge field theories is given in [18],[12]. In the case of 2D Yang-Mills theories (QCD_2), the path integral (45) can be explicitly done, which allows us to verify the method of reflected trajectories (formula (32)) by comparing its results with the known ones obtained in the framework of the loop approach [21],[22]. For QCD_2 , K is the Weyl cell (a polyhedron), S_F coincides with the affine Weyl group [12]. Contributions of trajectories reflected from the boundary ∂K (which is provided by the operator $\hat{Q}$ entering into the PI formula) are shown to be necessary to recover results of the loop approach [12].

The method can be also generalized to gauge models with fermions (with anticommuting (Grassmann) variables) [23],[6]. It is curious that the operator $\hat{Q}$ depends on Grassmann variables, which is important for a correct calculation of a semiclassical transition amplitude in minisuperspace cosmological gauge models with fermions [24].

References

- [1] R.P.Feynman and A.Hibbs, *Quantum Mechanics and Path Integrals* (Mc Graw-Hill, NY, 1965).
- [2] J.Klauder, *Path Integrals*, Acta Phys. Austrica Suppl, **XXII** (1980) 3.
- [3] W.Pauli, *Pauli Lectures on Physics* V.6 (MIT Press, 1973);
H.Kleinert and W.Janke, Lett.Nuovo Cim. **25** (1979) 297.
- [4] L.V.Prokhorov, Sov.J.Part.Nucl. **13**(5) (1982) 456.
- [5] H.Kleinert, *Path Intergals in Quantum Mechanics, Statistics, and Polymer Physics* (WSPC, Singapore, 1990).

- [6] L.V.Prokhorov and S.V.Shabanov, *Hamiltonian Mechanics of Gauge Systems* (JINR publication, B1-2-93-312, Dubna, 1993) (in Russian).
- [7] S.V.Shabanov, Phys.Lett.B **293** (1992) 117;
J.Phys.A: Math.Gen. **26** (1993) 2583.
- [8] P.A.M.Dirac, *Lectures on Quantum Mechanics* (Yeshiva Univ., N.Y., 1965).
- [9] L.V.Prokhorov, Sov.J.Nucl.Phys. **35** (1982) 192.
- [10] L.V.Prokhorov and S.V.Shabanov, Phys.Lett.B **216** (1989) 341.
- [11] L.V.Prokhorov and S.V.Shabanov, Sov.Phys.Uspekhi **34** (1991) 108.
- [12] S.V.Shabanov, *Phase space structure and the path integral for gauge theories on a cylinder*, Pisa University preprint, IFUP-TH 35/93 (Pisa, 1993), hep-th/9308002 (to appear in Phys.Lett.B).
- [13] J.E.Hetrick, *Canonical Quantization of Two Dimensional Gauge Fields*, preprint UvA-ITFA, Amsterdam, 1993, hep-th/9305020.
- [14] V.N.Gribov, Nucl.Phys.B **139** (1978) 1.
- [15] I.M.Singer, Commun.Math.Phys. **60** (1978) 7.
- [16] M.A.Soloviev, Theor.Math.Phys. (USSR) **78** (1989) 117.
- [17] S.V.Shabanov, *Lectures on Quantization of Gauge Theories by the Path Integral Method*, IFM preprint 16/92 (IFM, Lisbon, 1992) (to appear in “The Proceedings of the IV Hellenic School on High Energy Physics”, Corfu, Greece, 1992).
- [18] S.V.Shabanov, Phys.Lett.B **255** (1991) 398;
Mod.Phys.Lett.A **6** (1991) 909.
- [19] O.Bertolami, J.Mourao, R.F.Picken and I.P.Volobuev, Int.J.Mod.Phys.A **6** (1991) 4149.
- [20] S.V.Shabanov, Phys.Lett.B **272** (1991) 11.
- [21] A.A.Migdal, Sov.Phys.-JETP **42** (1976) 413.
- [22] G.Rajeev, Phys.Lett.B **212** (1988) 203.
- [23] S.V.Shabanov, J.Phys.A:Math.Gen. **24** (1991) 1199.
- [24] S.V.Shabanov, in: *Proceedings of the First Iberian Meeting on Gravity* (Evora, Portugal, September 1992) (to appear).